



Clinical Applications

- Cardiovascular Health Including Blood Pressure, Lipids and Clotting^{1,7}
- Reducing Inflammation of Joints, Skin, Other^{2,3}
- Reducing Back and Neck Pain⁴
- Support for Mental Health/Behavior⁵
- Support for the Reduction of Allergic-type Response⁶
- Glucose and Insulin Homeostasis^{7,8}

Omega 600 MAX™ and Pura Omega 780EC™ are among Vida Pura®'s exclusive line of pharmaceutical grade, enteric-coated, ultra-pure ArticOils® fish oils. These molecularly-distilled oils provide the respective milligrams of omega-3 essentials fatty acids per softgel, permitting dose flexibility. Independent third party assays document freshness and purity of every batch.

All Vida Pura® Formulas Meet or Exceed cGMP quality Standards

Discussion

At the same time as the media increases consumer awareness of the importance of fish oils, manufacturers face the challenge of eliminating heavy metals and PCBs. The industry-leading technologies used in the preparation of Vida Pura®'s Arctic Oils® surpass all national and international standards for environmental pollutants, including dioxins, PCBs, pesticides and heavy metals, including mercury.

Vida Pura®'s Omega oils are processed through special molecular distillation. Molecular distillation is a process commonly used to purify and concentrate fish oils. In this process, the fish oil is broken down into its basic molecular components and separated by molecular weight. Due to varying molecular weights, specific components can be removed or concentrated to the oil. This ensures that contaminants can be reduced to levels far below industry standards. It also allows the manufacturer to increase the concentration of active ingredients. For example, an 18:12 oil can be concentrated through molecular distillation to produce a 30:20 product as in the case of our Omega 600 MAX™.

Traditional molecular distillation technology separates contaminants from the oil through evaporation off a high temperature surface. Not all molecular distillation systems are the same. Below are further details about the systems employed for each of these ArticOils®.

Processing all three fish oil formulas begins with rigorous care and control of the starting raw materials (non-GMO sardines, mackerel, anchovies) to assure optimum oil quality. The oils are distilled in a controlled, pristine vacuum environment to minimize distillation temperatures. The exposure time even to these lower distillation temperatures is tightly controlled and uniform, resulting in levels of impurities well below the industry standards.

The technology employed for the Omega 600 MAX™ meets special Molecular Distillation Standards: As this oil is a concentrated source of Omega-3's, our manufacturer goes above and beyond the traditional purification methods to ensure its safety. This is accomplished by:

- A triple phase molecular distillation purification process to maximize purity while concentrating the EPA and DHA polyunsaturated fatty acids
- Ensuring consistency in contaminant removal and therefore purity levels through uniform processing times
- Reducing the evaporation stage to half the time of traditional systems to drive down the oxidative risk

The proprietary technologies used in the manufacture of Omega 600 MAX™ are in accordance with pharmaceutical standards that assure safe, consistent fish oils that exceed Council for Responsible Nutrition (CRN) Monograph standards and meet all international standards for oil safety including WHO, Prop 65, and more. They are third party tested according to AOCS (American Oil Chemical Scientists) international protocols to ensure purity and potency.⁹



The EPA and DHA present in fish oils can be concentrated either as free fatty acids, ethyl esters or reconstituted triglycerides. The debate on the comparative bioavailability of EPA and DHA as triglycerides versus ethyl esters is not yet settled. However, the physiological responses are similar when these fatty acids are administered over a week either as ethyl esters or triglycerides.¹⁰ The Omega 600 Max™/Pura Omega780 EC™ is the ethyl ester form, guaranteed to exceed the industry standards set by the CRN Monograph.

The International Society for the Study of Fats and Lipids made the following suggestions for dosing infants, children and lactating women in 2002 and updated the adult recommendation in 2004.¹¹

Category	Weight	Dose
Infants (1-18 months)	0-15 lbs	32 mg/lb EPA* +DHA
Children (1.5 - 15 yrs)		5 mg/lb EPA +DHA
Adults (15-100 yrs)		500 mg EPA +DHA
Lactating Women		300 mg DHA/day

It takes 2 gm/day of DHA supplementation for a month to saturate the plasma and three to six months of supplementation to saturate the tissues. Concentrations increase in breast milk within less than a week of DHA supplementation. Omega-3 fatty acids must be consumed regularly for maintenance of good health and prevention of disease.

Omega 600 MAX™

Supplement Facts		
Serving Size: 1 Softgel		
Servings Per Container: 120		
	Amount	%Daily Value*
Calories	12.7	
Calories from Fat	10.8	
Total Fat	1.2 g	2.1%
Vitamin E (as mixed tocopherols)	10 IU	33%
Natural Marine Lipid Concentrate	1.2 g	**
EPA (eicosapentaenoic acid)	360 mg	**
DHA (docosahexaenoic acid)	240 mg	**

* Percent Daily Value based on a 2,000 calorie diet.
** Daily value not established.
Other ingredients: Gelatin, glycerin, purified water, enteric coating and vanillin.

Pura Omega 780 EC™

Supplement Facts		
Serving Size: 1 Softgel		
Servings Per Container: 120		
	Amount Per Serving	%Daily Value*
Calories	13	
Calories from Fat	13	
Total Fat	1.4 g	3%
Vitamin E (as mixed tocopherols)	10 IU	33%
Natural Marine Lipid Concentrate	1.4 g	**
EPA (eicosapentaenoic acid)	450 mg	**
DHA (docosahexaenoic acid)	330 mg	**

* Percent Daily Value based on a 2,000 calorie diet.
** Daily Value not established.
Other ingredients: Gelatin, Glycerin, Purified Water, Enteric Coating (water, ethylcellulose, sodium alginate).

References

1. Calder PC. n-3 Fatty acids and cardiovascular disease: evidence explained and mechanisms expored. Clin Sci (Lond). 2004 Jul;107(1):1-11 [PMID: 15132735]
2. Storey A, McArdle F, Friendmann PS, Jackson MJ, Rhodes LE. Eicosapentaenoic acid and docosahexaenoic acid reduce UVB- and TNF alpha-induced IL-8 secretion in keratinocytes and UVB-induced IL-8 in fibroblasts. J Invest Dermatol. 2005 Jan;124(1):248-55. [PMID: 15654981]
3. Ki YJ, Kim HJ, No JK, Chung HY, Fernandes G. Anti-inflammatory action of dietary fish oil and calorie restriction. Life Sci 2006 Apr 18;78(21): 2523-32. Epub 2006 Jan 24 [PMID: 16438990]
4. Maroon JC, Bost JW. Omega-3 Fatty acids (fish oil) as an anti-inflammatory: an alternative to nonsteroidal anti-inflammatory drugs for discogenic pain. Surg Neurol. 2006 Apr; 65 (4):326-31 [PMID: 16531187]
5. Frangou S, Lewis M, McCrone P. Efficacy of ethyl-eicosapentaenoic acid in bipolar depression: randomized double-blind placebo-controlled study. Br J Psychiatry. 2006 Jan;188:46-50. [PMID: 16388069]
6. Kankaanpaa P, Sutas Y, Salminen S, Lichtenstein A, Isolauri E. Dietary fatty acids and allergy. Ann Med. 1999 Aug;31(4):282-7. [PMID: 10480759]
7. Ebbesson SO, Risica PM, Ebbesson LO, Kennish JM, Tejero ME. Omega-3 fatty acids improve glucose tolerance and components of the metabolic syndrome in Alaskan Eskimos: the Alaska Siberia project. Int J Circumpolar Health. 2005 Sep;64(4):396-408. [PMID:16277123]
8. Nettleton JA, Katz R. n-3 long-chain polyunsaturated fatty acids in type 2 diabetes: a review. J Am Diet Assoc. 2005 Mar;105(3):428-40 [PMID:15746832]
9. Molecular Distillation Reference Sheet. Bioriginal Food and Science Corp. Saskatchewan, CA
10. Hansen, J. B., J.O. Olsen, L. Wilsgard, V. Lyngmo and B. Svensson, Comparative effects of prolonged intake of highly purified fish oils as ethyl ester or triglyceride on lipids, haemostasis and platelet function in normolipaeic men. Eur J Clin Nutr 47, 497-507 (1993)
11. International Society for the Study of Fats and Lipids (ISSFAL, 2002)
12. Arterburn, LM. Dosing and distribution of n-3 fatty acids in humans. Presented at the Omega-3 fatty acids: Recommendations for therapeutics and prevention symposium. May 21, 2005. New York www.medscape.com/viewprogram/4605_pnt {accessed 6.9.06}

Precautions:

Avoid if allergic to any ingredient(s). Fish Oils may be associated with loose stools and abdominal discomfort in some patients. They may also increase the need for antioxidants. The Vida Pura Omega™ oils contain 10 IUs of vitamin E for added antioxidant protection.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.